



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$\left(\frac{x}{a} - \frac{y}{b}\right)^2 + \left(\frac{x}{a} + \frac{y}{b} + 1\right)\left(\frac{x}{a} + \frac{y}{b} - 1\right) = 0, \text{ i. e., } \frac{x^2}{a^2} + \frac{y^2}{b^2} = \frac{1}{2}.$$

Solved similarly by J. Scheffer.

256. Proposed by F. P. MATZ, Ph. D., Sc. D., Reading, Pa.

The bisectors of the four angles of any quadrilateral intersect in four points, all of which lie on the circumference of the same circle.

Solution by G. W. GREENWOOD, M. A., McKendree College, Lebanon, Ill.; FREDERIC R. HONEY, Ph. D., Trinity College, Hartford, Conn.; E. D. CARMICHAEL, Hartselle, Ala., and A. H. HOLMES, Brunswick, Me.

Denote the quadrilateral by  $ABCD$ , and the bisectors of the angles  $A, B, C, D$  by  $a, b, c, d$ , respectively. Suppose  $AB, DC$  intersect at  $O$ ;  $a, d$  at  $X$ ; and  $b, c$  at  $Y$ . We can easily show that, if  $B$  lies between  $A$  and  $O$ ,

$$\begin{aligned}\angle AXD &= \frac{1}{2}\pi + \frac{1}{2}\angle AOD; \\ \angle BYC &= \frac{1}{2}\pi - \frac{1}{2}\angle AOD.\end{aligned}$$

Hence, a pair of opposite angles of the quadrilateral whose sides are, consecutively,  $a, b, c, d$ , are supplementary, and it is therefore cyclic.

\* \* \* The problem admits of the following interesting extension: If  $a', b', c', d'$  are the bisectors, respectively, of the exterior angles  $A, B, C, D$ , the six quadrilaterals whose consecutive sides are, respectively,  $(abcd)$ ,  $(abc'd')$ ,  $(bcd'a')$ ,  $(cda'b')$ ,  $(dab'c')$ ,  $(a'b'c'd')$ , are cyclic. It is necessary that no two sides of the given figure be parallel.

GREENWOOD.

Also solved by F. P. Matz, J. Scheffer, G. B. M. Zerr, and W. J. Greenstreet. A. H. Holmes also solved No. 252.

## CALCULUS.

194. Proposed by G. W. GREENWOOD, M. A. (Oxon), Lebanon, Ill.

Show that the volume of the solid generated by the revolution of a segment of a circle, less than a semi-circle, about the diameter parallel to its chord, is equal to that of a sphere having a diameter equal to the chord; and hence that the volume is independent of the magnitude of the original circle, the length of the chord being known.

Solution by W. L. TRYON, Ithaca, N. Y.; E. D. CARMICHAEL, Hartselle, Ala.; and W. J. GREENSTREET, Editor of The Mathematical Gazette, Stroud, England.

Let the chord be of length  $2b$  in circle of radius  $a$ ; and take as axes the parallel diameter and the perpendicular bisector of the chord. The distance of the chord from the center is  $\sqrt{a^2 - b^2}$ . The required volume is

$$V = 2 \int_0^b (a^2 - x^2) dx - 2\pi b(a^2 - b^2) = 2\pi b(a^2 - \frac{1}{3}b^2) - 2\pi b(a^2 - b^2) = \frac{4}{3}\pi b^3.$$

This is independent of  $a$ , and is equal to the volume of a sphere of diameter  $2b$ .

Also solved by C. Hornung, J. Scheffer, A. H. Holmes, and the Proposer.